

Orthopaedic and Spine Institute
21 Spurs Lane, Suite 245, San Antonio, TX 78240
www.saspine.com
Tel# 210-487-7463

***POST-OP INSTRUCTIONS FOR PATIENTS WHO HAVE HAD
CLAVICLE FIXATION SURGERY***

Dressing: A dressing has been applied to your incision. The dressing may become moist from blood or drainage this is not a cause for alarm. However, if the drainage from the incision soaks through the dressing please call the office. This dressing may be removed 7 days after surgery, but please leave the adhesive strips in place over the incision. The incision may be left open to air, or, a small dressing applied.

Sling & Ice Cuff: The sling is worn for your comfort and protection. It should be taken off only as directed. The ice cuff is worn to minimize swelling and pain. It should be worn for the first 3 days, 3 hours on, 3 hours off, for comfort.

Wounds: Your wound should be kept clean and dry. You may shower once the dressing has been removed, unless otherwise directed by your physician.

Activity: Maintain the arm in the sling at all times (even in bed), except for three times daily to perform pendulum and elbow motion exercises and when showering, once the dressing has been removed. Do not place your arm above shoulder level. Strengthening might begin as early as six weeks after surgery if early healing is evident. No contact sports for at least 4-6 months.

Pain: Upon discharge from the hospital you should have a prescription for pain medication. Please take the pain medication with food. Follow the guidelines in the pain management protocol for

Orthopaedic and Spine Institute
21 Spurs Lane, Suite 245, San Antonio, TX 78240
www.saspine.com
Tel# 210-487-7463

using your medication or as directed by your physician. Do not drink alcoholic beverages or drive if you are using pain medications.

Precautions: If you develop a fever (temperature greater than 101 degrees Fahrenheit or 38.3 Celsius) or chills, or any undue symptoms such as unexpected pain, redness, swelling in your legs, rash, blisters, numbness, tingling, itching, hives or shortness of breath, please contact our office. If you have a problem or question, do not hesitate to call the office. The answering service will handle your calls to the office after hours and on weekends and one of the Spine and Orthopedic Institute clinicians will be available at 210-487-7463. **If you have an emergency, call 911 or go to the nearest emergency department.**

Diet: Start off with light meals and progress diet as tolerated. Carbohydrates will generally have a lower incidence of nausea and vomiting.

Follow Up: You will be seen 10-14 days after surgery for wound check and to begin a gentle range of motion program with physical therapy. Typically you will be seen again 6 weeks after surgery for xrays and then again at 12 weeks after surgery. Average time to complete healing after clavicle surgery is 16 weeks.